

I_VITAMIN_C_LEVELS_AND_ANTI OXIDANT_ACTIVITY_OF_PINEAP PLE_WET.pdf

by Rini Tri Hastuti G

Submission date: 02-Apr-2022 03:36AM (UTC+0700)

Submission ID: 1799178113

File name: I_VITAMIN_C_LEVELS_AND_ANTI-OXIDANT_ACTIVITY_OF_PINEAPPLE_WET.pdf (355.7K)

Word count: 4629

Character count: 23867

VITAMIN C LEVELS AND ANTIOXIDANT ACTIVITY OF PINEAPPLE WET CANDIED BASED ON THE LEVEL OF PINEAPPLES RIPENESS (*ANANAS COMOSUS* VAR. QUEEN) AS A FUNCTIONAL FOOD PRODUCT

RINI TRI HASTUTI^{1*}, PRADEA INDAH LUKITO²

^{1,2}Department of Pharmacy and Food Analysis, Health Polytechnic of Ministry of Health Surakarta, 57127, Indonesia

*Email: rini_triastuti@yahoo.com

39

Received: 10 Nov 2021, Revised and Accepted: 15 Dec 2021

ABSTRACT

Objective: The purpose of this study to find out the levels of vitamin C and antioxidant activity of pineapple wet candied based on the level of ripeness of pineapple as a functional food product.

Methods: This type of research was true experimental with posttest-only control research design. The determine vitamin C level method using spectrophotometry UV and the antioxidant activity testing using the DPPH (2,2-diphenyl-1-picryl-hydrazyl-hydrate) free radical inhibition method. Statistical test using One Way Anova test continued Post Hoc test.

Results: The results of the determination of vitamin C levels in pineapple wet candied with full green, half yellow and full yellow ripeness levels, respectively, were 4.22; 17.24; 0.75 mg/gram. The results of the antioxidant activity test (IC50) on wet candied pineapple with ripeness levels of full green, half yellow and full yellow, respectively, were 61.86; 49.98; and 72.11 mg/l. Based on the level of antioxidant intensity of each ripeness, the levels were strong, very strong and strong. The results of the One Way Anova test on the level of ripeness of pineapple fruit in wet candied for vitamin C levels and antioxidant activity both of them had a significant difference with P value = 0.000 (P<0.05) which indicates that there were a difference in antioxidant activity levels of wet candied pineapple based on the ripeness level of pineapple. The results of the Pos Hoc test showed a P value = 0.000 (P<0.05) with the highest antioxidant activity value on half-yellow pineapple wet candied.

Conclusion: The conclusion of this study is there are differences in vitamin C levels and antioxidant activity in pineapple wet candied based on the level of ripeness as a functional food product. Wet candied with half yellow pineapple contains the highest vitamin C and the highest antioxidant activity.

Keywords: Functional foods, Fruit ripeness level, Antioxidants, Vitamin C, Pineapple wet candied

© 2022 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<https://creativecommons.org/licenses/by/4.0/>)
DOI: <https://dx.doi.org/10.22159/ijap.2022.v14s1.06> Journal homepage: <https://innovareacademic.in/journals/index.php/ijap>

INTRODUCTION

Indonesia is a region with a tropical climate and a location in the equator. Indonesia allows the growth of various kinds of lush plants such as fruits and vegetables. Fruit contain various vitamins needed by the body, one of which is vitamin C acts as an antioxidant and is effective in overcoming free radicals that damage cells or tissues, including protecting the lens from oxidative damage caused by radiation [1].

Vitamin C is easily soluble in water; therefore when undergoing the process of slicing, washing and boiling foods containing vitamin C will experience a decrease in levels. The content of vitamin C in fruit and food will be damaged due to the oxidation process by the outside air, especially when heated. Storage is carried out at low temperatures (in the refrigerator) and cooking that does not cause color changes in foods containing vitamin C [2]. Vitamin C has anti-infective properties by helping to heal diseased or damaged body parts [3]. A person who is deficient in vitamin C is likely to reduce the antioxidant power in his body making it easy for stress, cancer sores, gingivitis and bone pain [4]. Vitamin C is useful for preventing cancer sores, maintaining healthy teeth and gums, and protecting from other diseases caused by vitamin C deficiency [5].

Pineapple is one type of fruit that is in great demand by the public. Pineapple is elliptical in shape, the skin of the fruit is scaly. The recommended vitamin C requirement is 30-60 mg per day, while the average vitamin C adequacy for a family is (53.7±2.2) mg [6]. Pineapple was known as a fruit rich of the enzyme bromelain, vitamin C, beta carotene and carotenoids. Vitamin C, beta carotene and carotenoids are known as free radical scavenging antioxidants, so consuming pineapple regularly can protect the body from various diseases, including cancer, coronary heart disease and premature aging [7]. Pineapple fruit in fresh condition only has a shelf life of 1-7 d at a temperature of ±22 °C [8]. The amount of pineapple production is

very abundant at harvest time, but it is not proportional to the level of consumption so that the selling price in the market is very cheap. Pineapple fruit is not utilized at the time of harvest, it is necessary to make efforts to extend the shelf life, increase the economic value and diversify the product so that it can be accepted by consumers [9].

The level of vitamin C in fruit is strongly influenced by the variety, environment, place of growth, the use of various types of fertilizers and the level of fruit maturity [10]. Classification of the level of ripeness of pineapples based on different skin colors from green to yellow to produce pineapples in 3 (three) categories, namely raw, ripe, and very ripe. Green pineapple fruit is included in the raw category, yellowish-green is included in the ripe category and full yellow is included in the very ripe category [11].

The results of the research by Syahrumsyah *et al.* (2010) showed that vitamin C levels in full green pineapple were 2.24%, half yellow pineapple were 1.92% and full yellow pineapple fruit was 1.75%. Raw fruits contain higher levels of vitamin C than ripe fruits. Vitamin C levels in fruit will increase until the fruit is ripe, and will decrease when the level of maturity has been exceeded. This decrease is due to the level of vitamin C in ripe fruit, which will turn into glucose [12]. Ripe fruit will increase water content, total dissolved solids, color, aroma, fruit texture, starch and sugar, while vitamin C content generally decreases [13].

Pineapple fruit that undergoes processing can increase shelf life and a wider marketing reach. This processing can be an added value to the income of pineapple farmers [14]. Pineapple fruit can be processed into various products such as pineapple jam, lunkhead, syrup, jelly, pineapple chips and Wet candied fruit. Processing pineapples into various kinds of products is one of the efforts so that pineapples are not wasted and rotten during the main harvest [10].

Wet candied is a preserved food with sugar and one form of processed food that is much liked by the public. The taste is sweet

Table 1: Characteristic of vitamin C in pineapple wet candied


Ripeness levels	Flavours	Colors	aroma	Texture
Full green	sweet, slightly sour	light yellow	Pineapple aroma	crunchy
Half yellow	sweet, slightly sour	light yellow	Pineapple aroma	crunchy
Full Yellow	sweet, slightly sour	yellow	Pineapple aroma	crunchy

Table 2: Identification of vitamin C in pineapple Wet candied with KMnO₄

Subject	Indication test	Result	Interpretation
Ascorbic acid	The purple color of KMnO ₄ disappears and becomes brown	The purple color of KMnO ₄ disappears and becomes brown	+
Destiluted water	disappears and becomes brown	The purple color of KMnO ₄ was stay	-
Full green pineapple wet candied		The purple color of KMnO ₄ disappears and becomes brown	+
Half yellow pineapple wet candied		The purple color of KMnO ₄ disappears and becomes brown	+
Full yellow pineapple wet candied		The purple color of KMnO ₄ disappears and becomes brown	+

Note: (+) = contains vitamin C, (-) = does not contain vitamin C

Identification of vitamin C with KMnO₄ reagent [23] occurred reduction and oxidation (redox) reactions. Vitamin C acts as a reducing agent, which is oxidized, while KMnO₄ acts as an oxidizing agent, which is reduced. Vitamin C can reduce permanganate ions because permanganate can be reduced in an acidic environment to manganese ions, while vitamin C is oxidized by permanganate ions because it has the potential to release H⁺ ions into dehydroascorbic acid. Ion permanganate accepts electron ions that escape from vitamin C so the purple color of KMnO₄ disappears [24]. The addition of potassium permanganate reagent to pineapple wet candied results in a changing color of KMnO₄ from purple to brown which indicates the formation of manganese ions (Mn²⁺). The reaction mechanism of vitamin C compounds with KMnO₄ reagent can be seen below:



Determination of vitamin C levels

Determination of vitamin C levels in pineapple wet candied was carried out using the UV-Vis spectrophotometry method because ascorbic acid is a compound that have chromophore and auxochrome groups in its structure so that it can be detected by UV detectors and provides absorption at UV wavelengths, in addition to UV spectrophotometry methods. UV-Vis has the advantages of being fast easy, has a low detection limit and has a high level of accuracy and precision [37]. [25]. In this research, the maximum wavelength is searched; the maximum wavelength is the wavelength that gives the largest absorption and is the same in each concentration. The maximum wavelength of a compound can be different if it is determined under different conditions and tools, so before determining the concentration, the maximum wavelength of ascorbic acid is determined. Determination of the maximum wavelength in this study was carried out by measuring the absorbance value of 10 ppm ascorbic acid solution in the wavelength range of 250-290 nm. The results of the absorbance measurement of 10 ppm ascorbic acid obtained the highest absorbance value at a wavelength of 266 nm with an absorbance value of 0.676.

Vitamin C levels [9] samples of pineapple wet candied can be determined using a calibration curve by measuring the absorbance of the sample and then calculated using a linear regression equation. The regression equation (fig. 2) generated from the calibration curve is $y = 0.0588x + 0.088$ and the correlation coefficient (R²) is 0.9985.

Based on the research results, it is known that the calibration curve obtained is linear because the value (R²) is close to 1, the correlation coefficient is used to determine the linearity of an analytical method. The correlation coefficient value close to 1 indicates the relationship between absorbance and concentration has a linear correlation where all points lie on a straight line [26]. Determination of vitamin C levels in candied pineapple fruit (table 3) was carried out using the UV-Vis spectrophotometry method, whose absorbance was measured at a wavelength of 266 nm.




Fig. 2: The calibration curve of vitamin C

Table 3: Calculation results of vitamin C levels in wet candied pineapple

Subject	Levels average±SD (mg/gram)
Full green pineapple wet candied	4.22±0.171
Half yellow pineapple wet candied	17.24±0.189
Full yellow pineapple wet candied	0.75±0.047

Note: The data were presented in mean±SD, n=3

The results of the calculation of the average vitamin C content of wet candied pineapple±SD value obtained indicate that the method used has a high precision value. The precision criterion is determined based on the value of the resulting standard deviation is small and close to zero, the smaller the SD value indicates, the more precise the resulting data [27]. According to Putri and Setiawati (2015) the recommended ADI (Nutritional Adequacy Rate) of vitamin C is 30-60 mg per day, while intake 2000 mg of half yellow pineapple wet candied it has been able to suitable the ADI, this is due to the processing process.

Based on the results of this study, it can be seen that half yellow pineapple wet candied has the highest average of vitamin C content, compared the candied full green, and full yellow pineapple candied. Based on the results of this study, it can be seen that vitamin C levels are influenced by the level of maturity, vitamin C levels in fruit will increase until the fruit is half ripe, and will decrease when the level of maturity overs, this is because vitamin C levels in overripe fruit will turns into glucose. Ripe fruit will increase water content, total dissolved solids, color, aroma, fruit texture, starch and sugar, while vitamin C content generally decreases [13].

Antioxidant activity test DPPH method

The antioxidant activity test of wet candied fruit with variations in the level of fruit maturity was carried out using the DPPH method

principle (table 4). The principle of measuring antioxidant activity using the DPPH reagent method is a change in the intensity of the purple color of DPPH, which is comparable with the concentration of the DPPH solution. DPPH free radicals which have unpaired electrons will give a purple color. Color will change turns yellow when the electrons are paired. The change in intensity of this purple color occurs due to the reduction of free radicals produced by the reaction DPPH molecule with a hydrogen atom. The hydrogen atom released by sample compound molecules to form 2,2-diphenyl-1-picrylhydrazine and causes DPPH color decay from purple to yellow. This color change will give a change in absorbance at length the maximum waveform of DPPH when measured using UV-Vis spectrophotometry [28].

Table 4: IC₅₀ value of antioxidant activity test

The levels of ripeness	Average±SD of IC ₅₀ (mg/l), n=3
Full green	61,86±0,75
Half yellow	49,98±0,85
Full yellow	72,11±0,69
Vitamin C	12,09±0,033

Note: The data were presented in mean±SD, n=3

Based on the antioxidant activity test on candied pineapple with various levels of ripeness of full green, half yellow, and full yellow, respectively, the average IC₅₀ value was 61,86; 49,98 and 72,11 mg/l (fig. 3).




Fig. 3: Antioxidant activity test result, Note: The data were presented in mean±SD, n=3

Based on the IC₅₀ value, it can be categorized as the potential antioxidant activity of pineapple wet candied with various levels of ripeness full green, half yellow and full yellow, the potential value respectively strong, very strong, and strong. The result of antioxidant activity can be influenced by the vitamin C content of each pineapple wet candied. The antioxidant activity of the three ripe pineapples on wet candied fruit was significantly different p value<0,05 (asym. sig=0,00. Based on the statistical test results, the antioxidant activity of vitamin C was significantly different (p value<0,05).

Vitamin C was a very strong antioxidant compound because it had a small IC₅₀ value, which is 12,09 g/ml (value is less than 50). Vitamin C have four hydroxyl groups. Vitamin C can react directly with the hydroxyl anion by donating one electron to form a semihydroascorbate compound which is not reactive and then undergoes a disproportionation reaction to form an unstable dehydroascorbate. Dehydroascorbate will be degraded to form oxalic acid and threonic acid. Vitamin C is a polar compound, the higher the concentration of vitamin C, the more abundant hydroxyl groups and the intensity of antioxidant activity increases [29].

CONCLUSION

The conclusion of this study is there are differences in vitamin C levels and antioxidant activity in pineapple wet candied based on the level of ripeness as a functional food product. Wet candied with half yellow pineapple contains the highest vitamin C and the highest antioxidant activity.

ACKNOWLEDGMENT

The researcher delivers the gratitude to Laboratory Management Unit of pharmacy and food analysis department, Health Polytechnic of Surakarta, for provided facilities for this research, so this research can be conducted well.

FUNDING

Personal charge

20

AUTHORS CONTRIBUTIONS

All the authors have contributed equally.

CONFLICT OF INTERESTS

The authors declare no conflict of interest

REFERENCES

- Rezvani HT, Moradi P, Soltani F. The effect of nitrogen fixation and phosphorus solvent bacteria on growth physiology and vitamin c content of Capsicum annum L. Iran J Plant Physiol. 2013;6:673-82.
- Biswas AK, Sahoo J, Chatli MK. A simple UV-vis spectrophotometric method for determination of β-carotene content in raw carrot, sweet potato and supplemented chicken meat nuggets. LWT- Food Sci Technol. 2011;44(8):1809-13. doi: 10.1016/j.lwt.2011.03.017.
- Rahman N, Ofika M, Said I. Analysis of vitamin C levels of gadung mango (Mangifera sp) and golek mango (Mangifera indica L.) based on maturity level using iodimetric method. J Agricul Food Chem J 2015;4:33-7.
- Gahler S, Otto K, Bohm V. Alterations of vitamin C, total phenolics, and antioxidant capacity as affected by processing tomatoes to different products. J Agricul Food Chem J 2003;51(27):7962-8. doi: 10.1021/jf034743q, PMID 14690380.
- Handrian RG, Meiriani H. Increasing levels of vitamin C in lowland tomato (Lycopersicon esculentum Mill.) with GA hormone. Agrotechn Onl J 2013;2:333-9.
- Putri MP, Setiawati YH. Analysis of vitamin C levels in fresh pineapple (Ananas comosus L.) and canned pineapple fruit with UV-vis spectrophotometry method. Wiyata J 2015;2:34-8.
- Lingga L. Hypertension free without drugs. Jakarta: Agro Media Library; 2012.
- Lies MS. Making various processed pineapple. Jakarta: Puspa Swara; 2001.

9. Syahrumsyah H, Murdianto W, Pramanti N. Effect of addition of carboxymethyl cellulose (CMC) and pineapple (*Ananas comosus* (L.) Merr.) maturity level on pineapple jam quality. *J Agricul Tech*. 2010;6:33-40.
10. Suyanti. Guidelines for processing 20 types of Fruit. Jakarta: Self-Help Spreader; 2010.
11. Lustini A, Saparudin PA. Classification of pineapple maturity levels using red-green-blue and hue-saturation-intensity color spaces. *Digit J Inf Tech*. 2019;2:22.
12. Oktoviana Y, Aminah S, Sakung J. Effect of storage time and sodium benzoate concentration on vitamin C levels of red chili (*Capsicum annum L*). *Chem Acad J* 2012;1:193-9.
13. Julianti E. Effect of maturity level and storage temperature on the quality of Dutch eggplant (*Cyphomandra betacea*). *Indonesian Hortic J* 2011;2:14-20.
14. Afrianti LH. 33 Kinds of fruits for health. Bandung: Alfabeta; 2010.
15. Muaris H, Fruit C. Gramedia Pu. 21 a Utama; 2003.
16. Puspita NS, Tamrin RA. Kajian Pembuatan minuman Fungsional dari daun Sirsak (*Annona muricata* Linn.) dengan penambahan Bubuk Jahe (*Zingiber officinale*). *J Sains Tekno Pangan*. 2019;4:2086-95.
17. Andriani Y. Ekstrak Betaadukan UAA. Dari *saccharomyces cerevisiae*. *J Gradien* 2007;27:26-30.
18. Hasnaeni H, Aminah A. Uji aktivitas antioksidan dan profil fitokimia ekstrak kayu beta-beta. *Lunasia Amara Blanco JFG* 2019;5(1):101-7. doi: 10.22487/j24428744.2019.v5.i1.12404.
19. Backer CA, Brink RCB. Flora of java. *Taxon*. 1966;15(2). doi: 10.1002/j.1365-8175.1966.tb00230.x.
20. Shaha RK, Rahman S, Asrul A. Bioactive compounds in chilli peppers (*Capsicum annum L.*) at various ripening (green, yellow and red) stages. *Annals Biol Res*. 2013;4:27-34.
21. Klimczak I, Malecka M, Szlachta M, Gliszczynska swiglo A. Effect of storage on the content of polyphenols, vitamin C and the antioxidant activity of orange juices. *J Food Compos Anal*. 2007;20(3-4):313-22. doi: 10.1016/j.jfca.2006.02.012.
22. Karinda M, Fatimawali, Citraningtyas CG. Comparison of results of determination of vitamin C levels of dodol mango using UV-vis spectrophotometry and iodometry methods. *J Pharmacol*. 2013;2:86-9.
23. Brand Williams W, Cuvelier EMME, Berset C. Use of a free radical method to evaluate antioxidant activity. *LWT-Food Sci Technol*. 1995;28(1):25-30. doi: 10.1016/S0023-6438(95)80008-5.
24. Chandra B Zulharmita, Putri WD. Determination of vitamin C and B1 levels in red dragon fruit (*Hylocereus lemairei* (Hook.) Britton and Rose) using UV-vis spectrophotometric method. *J Hygea Pharm*. 2019;11:62-74.
25. Ngibad K, Herawati D. Comparison of measurement of vitamin C levels using UV-vis spectrophotometry at UV and visible wavelengths. *Borneo J Medical Lab Technol*. 2019;1:77-81.
26. Selpiana E, Destiarti L, Nurlina. Comparison of Pb(II) determination methods in kapuas river by UV-vis spectrophotometry separate calibration method and standard addition. *J Equatorial Chem*. 2016;5:17-23.
27. Padmaningrum RT, Marwati S. Validation of cyclamate analysis methods by spectrophotometry and turbidimetry. *J Basic Sci*. 2015;4:23-9.
28. Molyneux P. The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity. *Songklan J Sci Technol*. 2004;26:211-21.
29. Lung JK, Destiani DP. Uji aktivitas antioksidan vitamin A, C, E dengan metode DPPH. *Farmaka*. 2017;15:53-62.

I_VITAMIN_C_LEVELS_AND_ANTIOXIDANT_ACTIVITY_OF_PINE...

ORIGINALITY REPORT

20%

SIMILARITY INDEX

15%

INTERNET SOURCES

13%

PUBLICATIONS

9%

STUDENT PAPERS

PRIMARY SOURCES

1	ppjp.ulm.ac.id Internet Source	1%
2	Submitted to School of Business and Management ITB Student Paper	1%
3	foodandnutritionresearch.net Internet Source	1%
4	www.foodandnutritionjournal.org Internet Source	1%
5	www.ijstr.org Internet Source	1%
6	Eva Martínez-Ispizua, Mary-Rus Martínez-Cuenca, José Ignacio Marsal, María José Díez et al. "Bioactive Compounds and Antioxidant Capacity of Valencian Pepper Landraces", <i>Molecules</i> , 2021 Publication	1%
7	Submitted to Republic Polytechnic Student Paper	1%

8	www.dovepress.com Internet Source	1 %
9	res.mdpi.com Internet Source	1 %
10	www.wjgnet.com Internet Source	1 %
11	"International meetings", Social Science & Medicine, 1983 Publication	1 %
12	journals.ums.ac.id Internet Source	1 %
13	Alexandra Stoll, Ricardo Salvatierra-Martínez, Máximo González, Jonathan Cisternas et al. "Importance of crop phenological stages for the efficient use of PGPR inoculants", Scientific Reports, 2021 Publication	1 %
14	sphinxsai.com Internet Source	1 %
15	garuda.ristekbrin.go.id Internet Source	1 %
16	www.coursehero.com Internet Source	<1 %
17	Annete de Jesus Boari Lima, Angelita Duarte Corrêa, Adelir Aparecida Saczk, Mariana	<1 %

Pereira Martins, Rachel Oliveira Castilho.
"Anthocyanins, pigment stability and
antioxidant activity in jaboticaba [*Myrciaria
cauliflora* (Mart.) O. Berg]", *Revista Brasileira
de Fruticultura*, 2011

Publication

18

T W Putri, N F Nursida, I Raya. "Antioxidant
Activity of *Chlorella vulgaris* Used As an
Antioxidant Cream", *Journal of Physics:
Conference Series*, 2021

Publication

<1 %

19

www.science.gov

Internet Source

<1 %

20

Submitted to University of Petra

Student Paper

<1 %

21

core.ac.uk

Internet Source

<1 %

22

Submitted to Universitas Sebelas Maret

Student Paper

<1 %

23

Submitted to University of Sheffield

Student Paper

<1 %

24

journal.unpad.ac.id

Internet Source

<1 %

25

Prayoga Byantara, Dianursanti. "Utilization of
Spirulina platensis microalgae as edible
coating to maintain quality of fresh

<1 %

strawberry (Fragaria sp.)", AIP Publishing,
2021

Publication

26

blog.longevity-and-antiaging-secrets.com

Internet Source

<1 %

27

jurnal.fkip.untad.ac.id

Internet Source

<1 %

28

www.scientific.net

Internet Source

<1 %

29

Sholichah Rohmani, Kezia Esther Dinda,
Ahmad Ainurofiq. "Formulation and
evaluation of the cream made from
potassium azeloyl diglycinate as an anti-
aging", Journal of Physics: Conference Series,
2021

Publication

<1 %

30

ejournal.sttif.ac.id

Internet Source

<1 %

31

injec.aipni-ainec.org

Internet Source

<1 %

32

Submitted to University of KwaZulu-Natal

Student Paper

<1 %

33

doaj.org

Internet Source

<1 %

34

dspace.ugal.ro

Internet Source

<1 %

35	jgpt.co.in Internet Source	<1 %
36	www.frontiersin.org Internet Source	<1 %
37	F M Fadhlillah, W Oktaviani, R Mariani. " Antioxidant activity of Ethanol Extract, - Hexane fraction, Ethyl Acetate fraction and Water fraction of Garut Orange Leaves (Blanco) ", Journal of Physics: Conference Series, 2019 Publication	<1 %
38	LEIBOFF, S. L.. "COLLOIDAL GOLD : A METHOD FOR THE PREPARATION OF COLLOIDAL GOLD FOR THE LANGE TEST ON CEREBROSPINAL FLUID", Archives of Dermatology and Syphilology, 1928. Publication	<1 %
39	Wen Zhao, Nan Huo, Liang Cui, Xiaoying Li, Z. Y. Ou. "Propagation of temporal mode multiplexed optical fields in fibers: influence of dispersion", Optics Express, 2021 Publication	<1 %
40	jurnal.unpad.ac.id Internet Source	<1 %
41	www.jim.unsyiah.ac.id Internet Source	<1 %

42 R Suseno, Surhaini, S L Rahmi, F Yanti. <1 %
"Characteristics and sensory properties of lemongrass, roselle, and ginger formulation herbal tea", IOP Conference Series: Earth and Environmental Science, 2022
Publication

43 Rahma Micho Widyanto, Lulu Safira, Nur Faadiyah Sofian, Syifa Aulia Mardhiyati et al. <1 %
"Studies on the Antioxidant and Cytotoxic Potentials of the Peel Extract of Dacryodes rostrata", BIO Web of Conferences, 2021
Publication

44 Seyhun Yurdugül. "Preservation of quinces by the combination of an edible coating material, Semperfresh, ascorbic acid and cold storage", European Food Research and Technology, 05/2005 <1 %
Publication

45 www.centralreaders.net <1 %
Internet Source

46 www.mdpi.com <1 %
Internet Source

47 Oliveira, Ana, Maria Helena Gomes, Elisabete M.C. Alexandre, Domingos P.F. Almeida, and Manuela E. Pintado. "Impact of pH on phytochemical profile of pasteurized peach <1 %

purée during storage", Journal of Agricultural and Food Chemistry

Publication

48

Tri Oktarina Safitri, Ali Napiah Nasution, Chrismis Novalinda Ginting, Ermi Girsang. "The Effectiveness of Red Spinach Extract Ointment (*Amaranthus tricolor* L.) against Healing Second Degree Burns in Wistar Rat", 2021 IEEE International Conference on Health, Instrumentation & Measurement, and Natural Sciences (InHeNce), 2021

Publication

<1 %

49

www.jurnal.unsyiah.ac.id

Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off